

**SEASONAL STATUS OF
VANCOUVER BIRDS**
June 1, 2013 Version 6
compiled by Rick Toochin
Comments? Contact E-Fauna BC


AREA COVERED


Greater Vancouver and Vicinity from the International Boundary (but including Point Roberts, Washington) North to 49° 35' N (including Furry Creek) West to the middle of the Strait of Georgia and Howe Sound (including Bowen Island) East to 122° 33' W (Bradner Road or 288 th St.) in (Abbotsford and Maple Ridge), but including all of Golden Ears Provincial Park.

Computer design and data entry by Rick Toochin

Key:	
█	Common: 25+ seen or heard daily
▬	Fairly common: 5-25 seen-heard daily
▬	Uncommon: 0-5 seen or heard daily
—	Rare: 1-10 records per year
.....	Casual: 2-20+ records; on average not seen every year; somewhat out of normal range
○	Accidental: single record of species; usually far outside normal range (1 - 4 days stay)
----->	Accidental: duration of stay (5+ days stay)
[I]	Introduced: species established for more than 10 years with a historically stable population
[X]	Extirpated: once a resident but no longer found in checklist area
*	Known to nest in checklist area
(#)	Refers to the number of documented sightings of a casual or accidental species in the checklist area; only 1 through 10 are shown
sr	Sight Record: a sighting by single or multiple observer(s) where there is only an acceptable written description but there is no physical evidence by way of a photograph, tape recording or skin specimen.
<i>Italicized species are not seen every year.</i>	

Total: 411 species, including 169 breeding species and 138 species seen less than once per year, 10 introduced and 9 extirpated species.





- Fulmars & Shearwaters**
 — *Northern Fulmar (3)*
 — *Sooty Shearwater (6) sr*
 — *Short-tailed Shearwater (2) sr*
 — *Black-vented Shearwater (1) sr*

- Storm-Petrels**
 — *Fork-tailed Storm-Petrel*
 — *Leach's Storm-Petrel (4)*


- Pelicans**
 — American White Pelican
 — Brown Pelican


- Cormorants**
 — Brandt's Cormorant
 — Double-crested Cormorant*
 — Pelagic Cormorant*


- Bitterns & Herons**
 — American Bittern*
 — *Least Bittern (1)*
 — Great Blue Heron*
 — Great Egret
 — *Snowy Egret (8)*
 — Cattle Egret
 — Green Heron*
 — Black-crowned Night-Heron*


- Vultures**
 — Turkey Vulture


- Ospreys, Kites, Eagles & Hawks**
 — Osprey*
 — *White-tailed Kite*












- Goatsuckers**
- *Lesser Nighthawk (1)*
- Common Nighthawk*
- *Common Poorwill (1)*


- Swifts**
- Black Swift
- Vaux's Swift*
- *White-throated Swift (6)*


- Hummingbirds**
- *Ruby-throated Hummingbird (3) sr*
- *Black-chinned Hummingbird (3) sr*
- Anna's Hummingbird*
- *Costa's Hummingbird (6)*
- Calliope Hummingbird
- Rufous Hummingbird*

- Kingfishers**
- Belted Kingfisher*

- Woodpeckers**
- Lewis's Woodpecker*
- *Acorn Woodpecker (1)*
- *Williamson's Sapsucker (1) sr*
- *Yellow-bellied Sapsucker*
- Red-naped Sapsucker
- Red-breasted Sapsucker*
- Downy Woodpecker*
- Hairy Woodpecker*
- American Three-toed Woodpecker*
- *Black-backed Woodpecker (1) sr*
- Northern Flicker*
- Pileated Woodpecker*







Wrens & Dipper

- *Rock Wren*
- Bewick's Wren*
- House Wren*
- Pacific Wren*
- *Sedge Wren (1)*
- Marsh Wren*
- American Dipper*

Kinglets & Gnatcatcher

- Golden-crowned Kinglet*
- Ruby-crowned Kinglet*
- *Blue-gray Gnatcatcher (4)*

Thrushes


- *Red-flanked Bluetail (1)*
- *Northern Wheatear (1)*
- *Western Bluebird*[X](5)*
- Mountain Bluebird*
- Townsend's Solitaire*
- *Veery*
- Swainson's Thrush*
- Hermit Thrush*
- *Dusky Thrush (1)*
- *Fieldfare (1)*
- American Robin*
- Varied Thrush*


Mimics


- Gray Catbird*
- Northern Mockingbird
- *Sage Thrasher (7)*
- *Brown Thrasher (4)*

Starlings

- European Starling* [I]
- *Crested Myna* [I,X]*








- Summer Finches**
 — *Rose-breasted Grosbeak*
 — Black-headed Grosbeak*
 — Lazuli Bunting*
 — *Indigo Bunting*
 — *Painted Bunting (2)*

- Blackbirds**
 — *Bobolink*
 — Red-winged Blackbird*
 — Western Meadowlark*
 — Yellow-headed Blackbird*
 — Rusty Blackbird
 — Brewer's Blackbird*
 — *Common Grackle (8)*
 — Brown-headed Cowbird*

- Orioles**
 — *Hooded Oriole (4)*
 — Bullock's Oriole*
 — *Baltimore Oriole (2)*

- Winter Finches**
 — *Brambling*
 — Gray-crowned Rosy-Finch
 — Pine Grosbeak
 — Purple Finch*
 — *Cassin's Finch*
 — House Finch*
 — Red Crossbill*
 — White-winged Crossbill
 — Common Redpoll
 — Pine Siskin*
 — *Lesser Goldfinch (3)*
 — American Goldfinch*
 — Evening Grosbeak*

- Old World Sparrows**
 — House Sparrow* [I]



Hypothetical List:

Hypothetical : species believed to have been correctly identified, but for which satisfactory documentation (a specimen, identifiable photograph, convincing field description) is lacking, at least in the season in question. In most cases the records listed here have not been reviewed.

Arctic Loon
Black-footed Albatross
Magnificent Frigatebird
White-faced Ibis
Swainson's Hawk
Mountain Plover
Little Curlew
Bristle-thighed Curlew
Long-toed Stint
American Woodcock
Lesser Black-backed Gull
Ross's Gull
Black-billed Cuckoo
Chimney Swift
Yellow-bellied Flycatcher
Phainopepla
Blue-headed Vireo
Golden-winged Warbler
Virginia's Warbler
Blackburnian Warbler
Mourning Warbler
Scarlet Tanager
Baird's Sparrow
Le Conte's Sparrow
Rustic Bunting
Hoary Redpoll